PhytoKeys 259: 199–215 (2025)
DOI: 10.3897/phytokeys.259.147785

Review Article

Aquatic macrophytes of Angola: a preliminary verified checklist

Rafael Somandjinga[1,2,3], Sílvia Quadros[1,2], Francisco Maiato[4,5,6], David Goyder[7]

1 *CIBIO, Research Centre in Biodiversity and Genetic Resources, InBio Associate Laboratory, BIOPOLIS Program in Genomics, Biodiversity and Land Planning; UNESCO Chair – Land Within Sea: Biodiversity & Sustainability in Atlantic Islands, University of the Azores, Rua da Mãe de Deus, 9500-321 Ponta Delgada, Portugal*
2 *Faculty of Science and Technology, University of the Azores, Azores, Portugal*
3 *Faculty of Natural Sciences, University of Namibe, Farol de Noronha Campus 274, Moçâmedes, Namibe, Angola*
4 *Mandume Ya Ndemufayo University, Avenida Hoji Ya Henda, 30, Lubango, Angola*
5 *Herbário do Lubango, ISCED-Huíla, Rua Sarmento Rodrigues S/N, Lubango, Angola*
6 *cE3c - Centre for Ecology, Evolution and Environmental Changes, Faculty of Sciences, University of Lisbon, Lisboa, Portugal*
7 *Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, London, UK*

Corresponding authors: Rafael Somandjinga (somandjinga2015@gmail.com); Sílvia Quadros (silvia.ab.quadros@uac.pt)

Abstract

Aquatic macrophytes are photosynthetic macroscopic organisms that grow permanently or periodically submerged or floating on the surface of the water. The aim of this study is to compile a checklist of fresh-water aquatic vascular plant macrophytes occurring in Angola, focusing on their origins, life forms, conservation status and distribution throughout the country. The checklist was compiled through a literature review of previous studies carried out in Angola, herbarium collections and online databases. A total of 526 species of macrophytes in 196 genera and 70 families was recorded. Cyperaceae was the most abundant family, followed by Poaceae, Lentibulariaceae, Lythraceae, Eriocaulaceae, Araceae, Podostemaceae, Hydrocharitaceae, Onagraceae and Plantaginaceae. The highest number of macrophytes was recorded from Huíla Province, followed by Moxico and Cuando Cubango Provinces, respectively, while Lunda Norte, Uíge and Zaire were poorly represented. From the aquatic macrophytes recorded, it was possible to identify the category of origin of 511 species. From these, 472 species are native, 23 introduced and 16 are endemic to Angola and the conservation status of these species was also assessed. The results obtained in this study show that there are still some gaps, especially with regard to the distribution of aquatic macrophytes in Angola, since many species are documented as native, but there is a lack of data on their distribution.

Key words: Angola, aquatic macrophytes, conservation status, distribution

OPEN ACCESS

Academic editor: Muthama Muasya
Received: 28 January 2025
Accepted: 21 May 2025
Published: 4 July 2025

Citation: Somandjinga R, Quadros S, Maiato F, Goyder D (2025) Aquatic macrophytes of Angola: a preliminary verified checklist. PhytoKeys 259: 199–215. https://doi.org/10.3897/phytokeys.259.147785

Copyright: © Rafael Somandjinga et al. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Aquatic macrophytes are macroscopic photosynthetic organisms, that grow permanently or periodically submerged or that float on the water surface (Chambers et al. 2008; Murphy et al. 2019; Kato et al. 2024). These plants have been recognised as a significant component of aquatic ecosystems, which play an important role in providing several environmental and ecosystem services (Sood et al. 2012; Vymazal 2013; Kumar et al. 2023); these

include, decontamination and recycling of nutrients in water (Khan et al. 2020; Kumar et al. 2020), control of water erosion (Kumar et al. 2023), biomass production (Amulya et al. 2023; Dębowski et al. 2023), act as organic fertilisers (Saleh 2023; Hernández-Fernández et al. 2024) and as water quality bio-indicators (Shelef et al. 2013; Bytyqi et al. 2020; Saleh 2023). In addition to these ecosystem services, aquatic macrophytes can also be used to perform wastewater treatment in extensive systems, such as constructed wetlands (Vymazal 2013).

Based on the habitat that these groups of plants occupy, macrophytes can be classified as: i) emergent macrophytes - rooted in submerged soils or land that are periodically flooded, with foliage that extends into the air; ii) floating leaf macrophytes - rooted in the bottom of lake or stream with leaves floating on the surface of the water; iii) free-floating macrophytes - plants floating on or under the surface of the water; iv) submerged macrophytes - plants that grows completely submerged under water (Chambers et al. 2008) and v) amphibious macrophytes - plants capable of living both in flooded areas and above water (Moura-Júnior and Cotarelli 2019).

Angola is a country with a rich biodiversity. However, vegetation studies in Angola are limited to individual or few published studies in scientific literature and most of the published literature and/or databases are not easily accessed (Huntley et al. 2019). The scarcity of vegetation studies of Angola was also mentioned by Figueiredo and Smith (2008), Figueiredo et al. (2009), Sosef et al. (2017), Goyder and Gonçalves (2019) and Goyder et al. (2023a). Although some studies have already been carried out, in most of these areas, the focus has been predominantly on the exploration of terrestrial groups (Figueiredo and Smith 2008; Figueiredo et al. 2009; Chisingui et al. 2018; Goyder et al. 2018; Gonçalves et al. 2021; Monteiro et al. 2022; Goyder et al. 2023a, 2023b; Almeida et al. 2024).

From the studies about aquatic macrophytes carried out in Angola, the collections of Hans Hess in early 1950 deserve special mention (Goyder and Gonçalves 2019). Other important work on aquatic macrophytes by Gomes (2009), who presented a vegetation study of the Angolan section of the Okavango Basin, referring to different aquatic macrophytes identified in this section of the Angolan territory, with about 23 species recorded. Goyder et al. (2018) presented a checklist of vascular plants, where they highlighted aquatic macrophytes of the Okavango Basin, stating that the wetlands of the region tend not to be botanically diverse and do not have local endemics. Therefore, although we verified some studies related to aquatic macrophytes, in Angola, there is no complete checklist of this important group of plants.

Compiling data into a checklist, based on fieldwork, bibliographic review, herbarium collections or online data platforms, allows us to increase knowledge related to biodiversity in a given study area, as well as to identify efficient tools to obtain scientific data (Moura-Júnior et al. 2013). This work aims to compile a preliminary verified checklist of the fresh-water aquatic vascular plant macrophytes in Angola, focusing on the category of origin, life form, conservation status and distribution throughout the country.

Material and methods

Study area

Angola is a country with an area of 1,246,700 square kilometres, located on the southwest coast of Africa, between latitude 4°22' and 18°02' South and longitude 11°41' and 24°05' East. It is bordered to the west by the Atlantic Ocean; to the north by the Republic of Congo and the Democratic Republic of Congo (DRC); to the east by the DRC and Zambia; to the south by Namibia (Huntley et al. 2019) (Fig. 1). According to the Koppen-Geiger climate classification, the northern region of Angola is typical of the tropical humid savannah group (Aw), the plateau is typical of the temperate mesothermal group (Cw), the southwest and the coastal plain are typical of the dry desert and semi-desert group (Bsh, Bwh) and the east has a temperate climate with hot summer rains (Cwa) (Huntley et al. 2019). Average annual rainfall varies from 37 mm in the desert province of Namibe to over 1600 mm in the provinces of Uíge, Lunda Norte and Lunda Sul. Angola is a country, rich in water resources; it contains nine major hydrological basins, six of which are transnational, serving as a “water tower” for much of southern and central Africa (Lourenco and Woodborne 2023); this Angolan water tower feeds major tributaries of the Congo, Zambezi and Okavango systems, as well as the Cuanza and Cunene Rivers (Milzow et al. 2009; Abiodun et al. 2019; Huntley et al. 2019; Lourenco et al. 2022).

Angola has been part of the Ramsar Convention since October 2021 (Ramsar 2025; Knight and Lourenco 2025) and, within the framework of the recommendations of this Convention, 11 candidate sites have been identified for inclusion on the list of Ramsar sites, namely: Lagunas do Mangal do Lobito (Benguela), Saco dos Flamingos (Luanda), Lagoa do Arco (Namibe), Parque Nacional de Cameia



Figure 1. Angola’s geographical location and mean annual precipitation.

(Moxico), Complexo das Zonas Húmidas da Lagoa do Carumbo (Lunda Norte), Lagoa do Calumbo (Luanda), Lagoa da Quilunda (Luanda) Complexo de Lagunas de Santiago Saurico (Bengo), Lagoa do Mangal do Chiloango (Cabinda), Baixo Cuanza (Luanda) and Complexo das Zonas Húmidas do Kumbilo Dirico (Cuando Cubango) (INBC 2021; Knight and Lourenco 2025). However, there is very little information on these wetlands and details are not yet available on the convention's website (Knight and Lourenco 2025). We also realise that, of these 11 wetlands, only two (Parque Nacional de Cameia (Moxico) and Complexo das Zonas Húmidas da Lagoa do Carumbo (Lunda Norte)) are located in the area with the highest average annual rainfall in the country, which indicates the scarcity of studies on wetlands.

In terms of its flora, Angola has around 6850 native plant species, a level of endemism of around 14.8% and 230 introduced species, four of which are considered highly invasive (Rejmánek et al. 2017; Goyder and Gonçalves 2019). According to Zigelski et al. (2019), fires, herbivory, extreme minimum temperatures and frost are factors that play significant roles in the floristic composition and physiognomic structure of the Angolan vegetation.

Collection of data

The checklist was drawn-up by conducting a literature review of previous botanical and floristic studies carried out in Angola (Costa et al. 2004; Figueiredo and Smith 2008; Gomes 2009; Goyder et al. 2018; Monteiro et al. 2022; Goyder et al. 2023b) and in the specialised literature in aquatic macrophytes (Fischer 1997; Germishuizen and Meyer 2003; Van Ginkel et al. 2011; Moura-Júnior et al. 2015; Cheek et al. 2017; Sieben et al. 2017; Ameka and Ernest 2019; Murphy et al. 2019; Pestana et al. 2024). Online databases, such as the International Union for the Conservation of Nature-Red list (IUCN 2024), Plants of the World Online (POWO 2024) and Global Biodiversity Information Facility (GBIF 2024) were also used. We also used the aquatic macrophyte collection of the Herbarium of Lubango (LUBA), housed at ISCED-Huíla, where we extracted the vouchers information from the specimens. Plant names were standardised by reference to POWO (2024) for Angiosperms, which incorporates the family concepts of APG IV (Angiosperm Phylogeny Group IV 2016) and Roux (2009) for Pteridophytes and Lycophytes. The species were classified according to life form, based on Gomes (2009), Moura-Júnior et al. (2015), Ameka and Ernest (2019), Moura-Júnior and Cotarelli (2019) and Pestana et al. (2024). The species were recorded by province, the codes for the Angolan provinces being based on Figueiredo and Smith (2008). For category of origin we used the terminology: native, endemic and introduced, also based on Figueiredo and Smith (2008) and POWO (2024). The conservation status of the species: Data Deficient (DD), Least Concern (LC), Near Threatened (NT), Vulnerable (VU), Endangered (EN) and Critically Endangered (CR) uses the IUCN (2024) categories and criteria.

Results

Diversity

A total of 526 species were registered, these belonging to 196 genera and 70 botanical families of aquatic macrophytes. The species were grouped according to major classification of plants: lycophytes (5 species), pteridophytes (20

species), dicots (176 species) and monocots (325 species) (Fig. 2). The botanical families with major number of species were Cyperaceae (155 spp.), Poaceae (70 spp.), Lentibulariaceae (21 spp.), Lythraceae (18 spp.), Eriocaulaceae (17 spp.), Podostemaceae (14 spp.), Araceae and Hydrocharitaceae (11 spp.) and Onagraceae and Plantaginaceae (10 spp.) (Table 1). These ten families together account for 60% of the total species recorded. Of all the species recorded in this study, 174 are deposited at LUBA.

The genera with the highest numbers of species are *Cyperus* L. (88), *Scleria* P.J.Bergius (21), *Utricularia* L. (16), *Eriocaulon* L. and *Rotala* L. (13) and *Panicum* L. (10) (Fig. 3). Its important to mention that these genera belong to the five richest families, mentioned above, namely: Cyperaceae (*Cyperus*, *Scleria*,), Lentibulariaceae (*Utricularia*), Eriocaulaceae (*Eriocaulon*), Lythraceae (*Rotala*) and Poaceae (*Panicum*). In contrast, 105 genera were represented by a single species.



Figure 2. Distribution of the number of species, genera and families for each category (Lycopodiophyta, Pteridophytes, Dicots and Monocots).

Table 1. Number of species (richness) distributed according to respective botanical family.

Family	Number of species
Cyperaceae	155
Poaceae	70
Lentibulariaceae	21
Lythraceae	18
Eriocaulaceae	17
Podostemaceae	14
Araceae, Hydrocharitaceae	11
Onagraceae, Plantaginaceae	10
Asteraceae, Fabaceae, Xyridaceae	9
Marsileaceae, Orchidaceae	8
Alismataceae, Commelinaceae, Polygonaceae	7
Droseraceae	6
Linderniaceae, Menyanthaceae, Nymphaeaceae, Pontederiaceae, Potamogetonaceae, Thelypteridaceae, Lamiaceae	5
Orobanchaceae, Araliaceae, Boraginaceae, Apocynaceae, Apiaceae	4
Aponogetonaceae, Convolvulaceae, Hydrostachyaceae, Juncaceae, Lycopodiaceae, Melastomaceae	3
Amaryllidaceae, Acanthaceae, Ceratophyllaceae, Gentianaceae, Haloragaceae, Iridaceae, Isoetaceae, Juncaginaceae, Dryopteridaceae, Ranunculaceae, Typhaceae, Rubiaceae, Parkeriaceae	2
Cannaceae, Amaranthaceae, Azollaceae, Brassicaceae, Cabombaceae, Celastraceae, Colchicaceae, Cymodoceaceae, Elatinaceae, Equisetaceae, Gleicheniaceae, Theophrastaceae, Ruppiaceae, Primulaceae, Ochnaceae, Mayacaceae, Marantaceae, Hypericaceae, Sphenocleaceae, Malvaceae	1



Figure 3. Distribution of species by genera.

Distribution of aquatic macrophytes by province

The distribution of macrophytes shows that Huíla Province is the best represented with a total 163 species, followed by Moxico and Cuando Cubango, Provinces with 74 and 71 species, respectively. Malanje (49 species), Cunene (43 species) and Namibe (41 species) come next (Fig. 4). The provinces with the least representation are: Lunda Norte, Uige and Zaire with fewer than 10 species. It should be noted that it was not possible to ascertain the distribution of 189 species.

Life forms

As far as biological forms are concerned, the species with the highest number of records were emergent macrophytes, with around 85%, followed by amphibious and floating-leaved macrophytes with around (6%) and finally free-floating and submerged macrophytes with around 2% (Fig. 5).

Category of origin of aquatic macrophytes

From the recorded aquatic macrophytes, it was possible to identify the category of origin of 511 species. From these, 472 species are native, corresponding to 91%, 23 especies are introduced, representing 4% and 16 are endemic to Angola, corresponding 3% (Fig. 6). The most representative, in terms of endemism are Cyperaceae (4 spp.), (*Cyperus gossweileri* Kük., *Cyperus subtenax* Kük., *Eleocharis cubangensis* H.E.Hess, *Scleria pulchella* Ridl.) and Lythraceae with three spp. (*Rotala nummularia* Welw. ex Hiern, *R. thymoides* Exell, *Dissotis rhinanthifolia* (Brenan) A.Fern. & R.Fern.). The Poaceae family with five spp. (*Arundo donax* L, *Coix lacryma-jobi* L., *Paspalum conjugatum* P.J. Bergius, *P. urvillei* Steud., *Polypogon viridis* (Gouan) Breistr) represents the largest number of introduced species. It was not possible to identify the category of origin of 12 species (*Berula erecta* (Huds.) Coville, *Genlisea glandulosissima* R.E.Fr., *Nymphoides aquatica* (J.F. Gmel.) Kuntze, *Nymphoides indica* (L.) Kuntze subsp. *occidentalis* A.Raynal, *Micrargeriella*



Figure 4. Distribution of aquatic macrophyte by province, showing the high representation of Huíla Province.



Figure 5. Distribution of species by life form: Em (Emergent), Am (Amphibious), FL (Floating-leaved), S (submersed), FF (Free floating).

aphylla R.E.Fr., *Aponogeton abyssinicus* Hochst. ex A.Rich., *Cyperus subtrigonus* (C.B.Clarke) Kük., *Fuirena leptostachya* Oliv. var. *nudiflora* K.Schum., *Hypolytrum heterophyllum* Boeckeler, *Brachycorythis congoensis* Kraenzl., *Disa ochrostachya* Rchb. f., *Potamogeton polygonifolius* Pourr.) and three are doubtful (*Myriophyllum spicatum* L., *Sphenoclea zeylanica* Gaertn., *Potamogeton richardii* Solms).



Figure 6. Category of origin of the aquatic macrophytes recorded from this study.

Conservation status

This study recorded a total of 386 species (73%) included in the IUCN Red List (IUCN 2024) from which 366 were considered Least Concern (LC), nine were Data Deficient (DD), (*Drosera elongata* Exell & J.R.Laundon, *Hydrostachys insignis* Mildbr. & Reimers, *Genlisea glandulosissima* R.E.Fr., *Rotala fontinalis* Hiern, *Nymphaea divaricata* Hutch., *N. sulphurea* Gilg, *Eleocharis cubangensis* H.E.Hess, *Eriocaulon longipetalum* Rendle, *Xyris imitatrix* Malme.), six were Vulnerable (VU) (*Utricularia inflexa* Forssk., *U. stellaris* L.f., *Rotala smithii* A.Fern. & Diniz, *Inversodicraea cristata* Engl., *I. warmingiana* (Gilg) Engl. and *Ledermanniella aloides* (Engl.) C.Cusset), two Near Threatened (NT) (*Ammannia baccifera* L. and *Anagallis kochii* H.E.Hess), two Endangered (EN) (*Genlisea angolensis* R.D.Good and *Leiothylax quangensis* (Engl.) Warm.) and one Critically Endangered (CR) (*Scleria pulchella* Ridl.). The family Podostemaceae was the family with major number of species with Vulnerable status (VU), namely: *Inversodicraea cristata* Engl., *I. warmingiana* (Gilg) Engl. and *Ledermanniella aloides* (Engl.) C.Cusset. A total of 139 species are not listed on the IUCN Red List.

Discussion

Diversity

Globally, 3499 species of aquatic macrophytes are known (Lobato-de Magalhães et al. 2023; Lobato-de Magalhães et al. 2024), distributed in six main ecozones (Nearctic, Neotropics, Australasia, Palaearctic, Afrotropics, Orient). Angola is part of the Afrotropics ecozone, which includes a total of 916 species according to Lobato-de Magalhães et al. (2024). In this study, a total of 526 species of aquatic macrophytes were recorded throughout the country, representing aproximatelly 8% of a total of ca. 6850 plant species actually known from Angola (Goyder and Gonçalves 2019).

The total number of aquatic macrophytes recorded, represented mainly by Cyperaceae and Poaceae, were also reported in various studies (Moura-Júnior et al. 2015; Adams et al. 2016; Sieben et al. 2017; Ameka and Ernest 2019; Murphy et al. 2019; Córdova et al. 2022; Pestana et al. 2024). The success of these species in aquatic environments is maybe due to several factors, namely: rapid vegetative and sexual reproduction, which leads to a rapid increase in the population, the ability to regenerate from fragments, high phenotypic plasticity, as well as efficient dispersal mechanisms (Hill et al. 2020).

Distribution of aquatic macrophytes

Based on Angola's varied climatic zones, the various habitat types and the number of macrophytes distributed throughout the country, we can predict that the total number of species recorded is far from the potential we might expect from this vast territory. It should be emphasised that the provinces with the highest number of species in this study are not necessarily the richest and most diverse as shown in Fig. 4. This is because these provinces were more comprehensively documented botanically compared to others, due to many historical reasons and existence of research infrastructures, a colonial legacy, emphasising the need for more and continued studies in other unexplored areas of the country, especially in the provinces located in northern and eastern parts of the country. According to Sieben et al. (2021), there is a strong correlation between the relative number of species and the average annual rainfall. Therefore, these areas represent the Angolan provinces characterised by the highest average annual rainfall (> 1600 mm/yr), so it is to be expected that the number of aquatic macrophytes can be higher than that mentioned in the scientific literature. Another curious fact is related to Namibe Province, characterised with average rainfall (< 600/yr), being occupied mostly by the Namib Desert. According to these characteristics, a low number of aquatic macrophytes would be expected. However, this Province has a considerable number (41) of species. This study was based on the list presented by Figueiredo and Smith (2008), where many documented species do not show their distribution throughout the country and online databases also lack this information. Therefore, given this scenario, the disparity in terms of macrophyte distribution compared to other parts of the country is essentially related to greater effort in floristic surveys in southern parts, when compared to others and, on the other hand, due to the lack of information in databases about the distribution of many species of aquatic macrophytes in Angola.

Life form

The most predominant life forms of macrophytes in our study were the emergent species, which can be explained by the ability of these plants to colonise a wide range of aquatic environments, from the beds themselves to the banks and to withstand long periods of drought (Hill et al. 2020). The correct recording of life forms is a fundamental aspect in the study of aquatic macrophytes (Moura-Júnior and Cotarelli 2019). However, this task has often become a major challenge since, in many cases, the labels on collections, herbarium records and some online platforms do not specify or adequately describe the life form and/or habitat, making it difficult to characterise this group of plants (Córdova et al. 2022). Another point that deserves special attention is the inclusion of amphibian species in aquatic macrophyte inventories, since, in some cases, errors can be made due to difficulties in recognising the physical boundary between wet and dry areas. Additionally, on the other hand, due to environmental changes that can occur days before sampling in the field, such as the rapid rise in water levels in an ecosystem due to heavy rainfall (Moura-Júnior and Cotarelli 2019).

Category of origin of aquatic macrophytes

In terms of origin categories, the number of endemic species recorded was 16, which represents 3% of total number of species recorded. However, this number may be higher, since classification doubts arose with macrophytes of some endemic genera. It is worth mentioning here some species that are referred to in literature as invasive in various parts of the world and which are recorded in this study, namely *Pontederia crassipes* Mart. (Datta et al. 2021; Ghoussein et al. 2023), *Pistia stratiotes* L. (Ameka and Ernest 2019; Hill et al. 2020) *Mimosa pigra* L. (Shanungu 2009; Welgama et al. 2022), *Cyperus papyrus* L. (Ameka and Ernest 2019), *Vossia cuspidata* Griff. (Ameka and Ernest 2019; Mahmoud et al. 2021), *Typha domingensis* Persoon (Edegbene 2018; Ameka and Ernest 2019) *Phragmites australis* (Cav.) Steud (Rohal et al. 2019; Coleman et al. 2023), *Arundo donax* L. (Hardion et al. 2014; Hardesty-Moore et al. 2020; Jiménez-Ruiz et al. 2021; Míguez et al. 2022). A study on invasive plants in Angola, conducted by Rejmánek et al. (2017), points to two of the invasive species listed above (*Pontederia crassipes* Mart. and *Arundo donax* L.), the latter of which has been spotted in five of the 14 vegetation types in western Angola, more than once and in dense stands. Since these plants cause disturbances in aquatic ecosystems, research should be carried out to understand the behavioural patterns of these species in Angola and, therefore, draw up control and monitoring programmes to guarantee the recovery of the targeted ecosystems.

Conservation status of species

The vast majority (95%) of the species assessed were considered to be of Least Concern. However, four species stand out here: one listed as Critical Endangered (CR) (*Scleria pulchella* Ridl.) which, according to Larridon and Bauters (2020), has only been recorded three times, the last being in 1952, near the Humpata and Palanca Rivers in Huíla Province. Three others classified as Endangered (EN) are: *Saxicolella angola* Cheek which was classified as endangered by Cheek et al. (2022) according to IUCN category and criterion B2ab(iii) (2012), *Genlisea angolensis* R.D.Good considered as a rare species, listed as endangered due to various current threats that are causing a decrease in its habitat and *Leiothylax quangensis* (Engl.) Warm. which can be found in Cameroon, Angola and the Democratic Republic of Congo (DRC); however, its distribution in Angola is unknown (Ghogue 2010a; Ghogue 2010b). It is worth mentioning a species of the Podostemaceae family (*Angolaea fluitans* Wedd.) which, although not formally evaluated, Goyder and Gonçalves (2019) suggest it may be extinct due to the construction of hydroelectric dams along the Cuanza River.

The species considered vulnerable include: *Rotala smithii* A.Fern. & Diniz known only from the DRC and Angola. It has been recorded in only two locations in the DRC and a similar distribution is expected to occur also in Angola. This species is threatened by water pollution (Ghogue 2010c). *Inversodicraea cristata* Engl. is known from only a few sites and occupies a much specialised habitat; some of these sites are threatened by habitat disturbance and degradation due to trampling by tourists (Ghogue 2017). *Inversodicraea warmingiana* (Gilg) Engl. is endemic to southern Angola, in the Cubango River, which then becomes the

Okavango River and northern Namibia. Its population is in decline due to environmental degradation (Sieben 2018). *Ledermanniella aloides* (Engl.) C.Cusset has been recorded globally in only five locations and Angola in only one (Diop 2010). The fact that 140 out of 526 species of aquatic macrophytes are not included on the IUCN Red List is worrying. Research in this area should be encouraged.

Conclusions

This study sets a precedent in research into the diversity, distribution, life forms and conservation status of aquatic macrophytes in Angola. Our results show that the Cyperaceae and Poaceae stand out as the most diverse and, in terms of life forms, the category of emergent macrophytes is the most dominant. As for the representativeness of aquatic macrophytes throughout the country, there is an uneven distribution, with the southern provinces being the best documented as a result of the larger floristic surveys that have been carried out.

Native species are the most prevalent, but the presence of introduced and invasive species requires investigation in order to understand the behaviour patterns of these species in Angola and the implementation of continuous monitoring programmes in order to preserve native ecosystems.

The results obtained in this study show that there are still some gaps, especially with regard to the distribution of aquatic macrophytes in Angola, since many species are documented as native, but there is a lack of data on their distribution. In view of the above, we recommend continuing studies on the distribution patterns of aquatic macrophytes in Angola and investing in floristic survey studies in the country's various provinces, in order to fill in the gaps in knowledge and, thus, promote the preservation of Angola's aquatic ecosystems.

Acknowledgements

We are grateful to the Herbarium LUBA team, in particular José João Tchamba, for giving us access to their aquatic macrophyte collection database. We are also grateful to Bernardino Bambi for preparing the maps for this work. Funding for open access charge: CIBIO - Açores.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by FCT - Fundação para a Ciência e Tecnologia, I.P. with the reference PRT/BD/154838/2023.

Author contributions

Writing – original draft: RS. Investigation: RS, SQ, FM, DG. Writing – review and editing: RS, SQ, FM, DG. Supervision: SQ, FM. Data curation: SQ, FM, DG.

Author ORCIDs

Rafael Somandjinga https://orcid.org/0009-0005-4852-7703
Sílvia Quadros https://orcid.org/0000-0001-8562-6640
Francisco Maiato https://orcid.org/0000-0002-8859-7491
David Goyder https://orcid.org/0000-0002-3449-7313

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

Abiodun BJ, Makhanya N, Petja B, Abatan AA, Oguntunde PG (2019) Future projection of droughts over major river basins in Southern Africa at specific global warming levels. Theoretical and Applied Climatology 137: 1785–1799. https://doi.org/10.1007/s00704-018-2693-0

Adams JB, Veldkornet D, Tabot P (2016) Distribution of macrophyte species and habitats in South African estuaries. South African Journal of Botany 107: 5–11. https://doi.org/10.1016/j.sajb.2016.08.001

Almeida RF, Antonio-Domingues H, Maiato PGF, Goyder DJ (2024) A new Angolan species from the Triaspis hypericoides complex (Malpighiaceae) based on macromorphology and palynology. Nordic Journal of Botany 2024(7): e04336. https://doi.org/10.1111/njb.04336

Ameka GK, Ernest A (2019) Aquatic macrophyte diversity, distribution, and control in coastal Southern Ghana, with priority on non-native invasive species. Indo Pacific Journal of Ocean Life 3(2): 87-93. https://doi.org/10.13057/oceanlife/o030205

Amulya K, Morris S, Lens PN (2023) Aquatic biomass as sustainable feedstock for biorefineries. Biofuels, Bioproducts & Biorefining 17(4): 1012–1029. https://doi.org/10.1002/bbb.2471

Angiosperm Phylogeny Group IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. Botanical Journal of the Linnean Society 181: 1–20. https://doi.org/10.1111/boj.12385

Bytyqi P, Czikkely M, Shala-Abazi A, Fetoshi O, Ismaili M, Hyseni-Spahiu M, Millaku F (2020) Macrophytes as biological indicators of organic pollution in the Lepenci River Basin in Kosovo. Journal of Freshwater Ecology 35(1): 105–121. https://doi.org/10.1080/02705060.2020.1745913

Chambers PA, Lacoul P, Murphy KJ, Thomaz SM (2008) Global diversity of aquatic macrophytes in freshwater. Hydrobiologia 595: 9–26. https://doi.org/10.1007/s10750-007-9154-6

Cheek M, Feika A, Lebbie A, Goyder DJ, Tchiengue B, Sene O, Van der Burgt X (2017) A synoptic revision of Inversodicraea (Podostemaceae). Blumea 62(2): 125–156. https://doi.org/10.3767/blumea.2017.62.02.07

Cheek M, Molmou D, Magassouba S, Ghogue JP (2022) Taxonomic revision of Saxicolella (Podostemaceae), African waterfall plants highly threatened by Hydro-Electric projects. Kew Bulletin 77(2): 403–433. https://doi.org/10.1007/s12225-022-10019-2

Chisingui AV, Gonçalves FM, Tchamba JJ, Luis JC, Rafael MFF, Alexandre JL (2018) Vegetation survey of the woodlands of Huíla Province. Biodiversity & Ecology 6: 426–437. https://doi.org/10.7809/b-e.00355

Coleman DJ, Cassalho F, Miesse TW, Ferreira CM (2023) The role of invasive Phragmites australis in wave attenuation in the Eastern United States. Estuaries and Coasts 46(2): 404–416. https://doi.org/10.1007/s12237-022-01138-x

Córdova MO, Keffer JF, Giacoppini DR, Pott VJ, Pott A, Moura-Júnior EG, Munhoz CBR (2022) Aquatic macrophytes in southern Amazonia, Brazil: Richness, endemism, and comparative floristics. Wetlands 42(4): 27. https://doi.org/10.1007/s13157-022-01545-7

Costa E, Martins T, Monteiro F (2004) A checklist of Angola grasses-Checklist das Poaceae de Angola. Southern African Botanical Diversity Network Report No. 28. SABONET, Pretoria.

Datta A, Maharaj S, Prabhu GN, Bhowmik D, Marino A, Akbari V, Kleczkowski A (2021) Monitoring the spread of water hyacinth (Pontederia crassipes): Challenges and future developments. Frontiers in Ecology and Evolution 9: 631338. https://doi.org/10.3389/fevo.2021.631338

Dębowski M, Zieliński M, Kazimierowicz J, Walery M (2023) Aquatic Macrophyte Biomass Periodically Harvested Form Shipping Routes and Drainage Systems in a Selected Region of Poland as a Substrate for Biogas Production. Applied Sciences 13(7): 4184. https://doi.org/10.3390/app13074184

Diop FN (2010) *Ledermanniella aloides*. The IUCN Red List of Threatened Species 2010: e.T185425A8409077. https://doi.org/10.2305/IUCN.UK.2010-3.RLTS.T185425A8409077.en

Edegbene AO (2018) Invasive grass (Typha domingensis): A potential menace on the assemblage and abundance of migratory/water related birds in Hadejia-Nguru wetlands, Yobe State, Nigeria. Tropical Freshwater Biology 27(2): 13–30. https://doi.org/10.4314/tfb.v27i2.2

Figueiredo E, Smith GF (2008) Plants of Angola/plantas de Angola. –. Strelitzia 22: 279.

Figueiredo E, Smith GF, César J (2009) The flora of Angola: First record of diversity and endemism. Taxon 58: 233–236. https://doi.org/10.1002/tax.581022

Fischer E (1997) A revision of the gem Dopatrium (Scrophulariaceae-Gratioloideae). -. Nordic Journal of Botany 17: 527–555. https://doi.org/10.1111/j.1756-1051.1997.tb00349.x

GBIF (2024) Global Biodiversity Information Facility. www.gbif.org [Accessed: 28 October 2024]

Germishuizen G, Meyer NL [Eds] (2003) Plants of southern Africa: an annotated checklist. Strelitzia 14. National Botanical Institute, Pretoria.

Ghogue JP (2010a) *Genlisea angolensis*. The IUCN Red List of Threatened Species 2010: e.T185240A8373311. https://doi.org/10.2305/IUCN.UK.2010-3.RLTS.T185240A8373311.en

Ghogue JP (2010b) *Leiothylax quangensis*. The IUCN Red List of Threatened Species 2010: e.T185346A8392899. https://doi.org/10.2305/IUCN.UK.2010-3.RLTS.T185346A8392899.en

Ghogue JP (2010c) *Rotala smithii*. The IUCN Red List of Threatened Species 2010: e.T185345A8392697. https://doi.org/10.2305/IUCN.UK.20103.RLTS.T185345A8392697.en

Ghogue JP (2017) *Inversodicraea cristata*. The IUCN Red List of Threatened Species 2017: e.T185309A120459889. https://doi.org/10.2305/IUCN.UK.2017-3.RLTS.T185309A120459889.en

Ghoussein Y, Abou-Hamdan H, Fadel A, Coudreuse J, Nicolas H, Faour G, Haury J (2023) Biology and ecology of Pontederia crassipes in a Mediterranean river in Lebanon. Aquatic Botany 188: 103681. https://doi.org/10.1016/j.aquabot.2023.103681

Gomes AL (2009) Análise Técnica, Biofísica e Sócio Económica do lado Angolano da Bacia Hidrográfica do Rio Cubango: relatorio final Vegetação da parte Angolana da Bacia Hidrográfica do Rio Cubango.

Gonçalves FMP, Chisingui AV, Luís JC, Rafael MFF, Tchamba JJ, Cachissapa MJ, Caluvino IMC, Bambi BR, Alexandre JLM, Chissingui MDG, Manuel SKA, Jacinto HD, Finckh M, Meller P, Jürgens N, Revermann R (2021) First vegetation-plot database of woody species from Huíla Province, SW Angola. Vegetation Classification and Survey. 2: 109–116. https://doi.org/10.3897/VCS/2021/68916

Goyder DJ, Gonçalves FMP (2019) The flora of Angola: collectors, richness and endemism. In: Huntley BJ, Russo V, Lages F, Ferrand N (Eds) Biodiversity of Angola, science and conservation: a modern synthesis. Springer, 79–96. https://doi.org/10.1007/978-3-030-03083-4_5

Goyder DJ, Barker N, Bester SP, Frisby A, Janks M, Gonçalves FMP (2018) The Cuito catchment of the Okavango system: A vascular plant checklist for the Angolan headwaters. PhytoKeys 113: 1–31. https://doi.org/10.3897/phytokeys.113.30439

Goyder DJ, Davies N, Finckh M, Gomes A, Gonçalves FMP, Meller P, Paton AJ (2023a) New species of Asclepias (Apocynaceae), Baphia (Leguminosae), Cochlospermum (Bixaceae) and Endostemon (Lamiaceae) from the Kalahari sands of Angola and NW Zambia, with one new combination in Vangueria (Rubiaceae). PhytoKeys 232: 145. https://doi.org/10.3897/phytokeys.232.110110

Goyder DJ, Gomes AL, Gonçalves FMP, Luís JC, Darbyshire I (2023b) A botanical assessment of Mt Namba, Cuanza-Sul, Angola: An isolated mountain towards the northwestern limits of the Great Escarpment of southern Africa. Namibian Journal of Environment 8: 77–92.

Hardesty-Moore M, Orr D, McCauley DJ (2020) Invasive plant Arundo donax alters habitat use by carnivores. Biological Invasions 22(6): 1983–1995. https://doi.org/10.1007/s10530-020-02234-4

Hardion L, Verlaque R, Saltonstall K, Leriche A, Vila B (2014) Origin of the invasive Arundo donax (Poaceae): A trans-Asian expedition in herbaria. Annals of Botany 114(3): 455–462. https://doi.org/10.1093/aob/mcu143

Hernández-Fernández L, Vázquez JG, Hernández L, Campbell R, Martínez J, Hajari E, Lorenzo JC (2024) Use of Euclidean distance to evaluate Pistia stratiotes and Eichhornia crassipes as organic fertilizer amendments in Capsicum annuum. Acta Physiologiae Plantarum 46(2): 21. https://doi.org/10.1007/s11738-024-03652-w

Hill MP, Coetzee JA, Martin GD, Smith R, Strange EF (2020) Invasive alien aquatic plants in South African freshwater ecosystems. Biological Invasions in South Africa. Springer International Publishing, Cham, 97–114. https://doi.org/10.1007/978-3-030-32394-3_4

Huntley BJ, Beja P, Vaz-Pinto P, Russo V, Veríssimo L, Morais M (2019) Biodiversity conservation: history, protected areas and hotspots. In: Huntley BJ, Russo V, Lages F, Ferrand N (Eds) Biodiversity of Angola, science and conservation: a modern synthesis. Springer, 495–51. https://doi.org/10.1007/978-3-030-03083-4

INBC (2021) Sítios Ramsar de Angola: Zonas Húmidas de Importância Internacional – Vol. I.

IUCN (2024) The IUCN Red List of Threatened Species, Version 2024-1. https://www.iucnredlist.org/ [Accessed: 30 July 2024]

Jiménez-Ruiz J, Hardion L, Del Monte JP, Vila B, Santín-Montanyá MI (2021) Monographs on invasive plants in Europe N° 4: Arundo donax L. Botany Letters 168(1): 131–151. https://doi.org/10.1080/23818107.2020.1864470

Kato S, Shiga T, Yamanouchi T, Nishihiro J (2024) Database of aquatic macrophytes in Japanese lakes. Ecological Research 39(3): 368–379. https://doi.org/10.1111/1440-1703.12449

Khan S, Nawab J, Waqas M (2020) Constructed wetlands: a clean-green technology for degradation and detoxification of industrial wastewaters. Bioremediation of Industrial Waste for Environmental Safety: Vol. II: Biological Agents and Methods for Industrial Waste Management, 127–163. https://doi.org/10.1007/978-981-13-3426-9_6

Knight J, Lourenco M (2025) Wetlands of Angola: Locations, Biodiversity, and Conservation. Wetlands of Tropical and Subtropical Asia and Africa: Biodiversity, Livelihoods and Conservation, 137–160. https://doi.org/10.1002/9781394235278.ch7

Kumar V, Kumar P, Singh J, Kumar P (2020) Potential of water fern (*Azolla pinnata* R. Br.) in phytoremediation of integrated industrial effluent of SIIDCUL, Haridwar, India: Removal of physicochemical and heavy metal pollutants. International Journal of Phytoremediation 22(4): 392–403. https://doi.org/10.1080/15226514.2019.1667950

Kumar S, Singh R, Kumar D, Bauddh K, Kumar N, Kumar R (2023) An Introduction to the Functions and Ecosystem Services Associated with Aquatic Macrophytes. In Aquatic Macrophytes: Ecology, Functions and Services. Springer, Singapore, 1–20. https://doi.org/10.1007/978-981-99-3822-3_1

Larridon I, Bauters K (2020) *Scleria pulchella*. The IUCN Red List of Threatened Species 2020: e.T120180747A172033420. https://doi.org/10.2305/IUCN.UK.2020-3.RLTS.T120180747A172033420.en

Lobato-de Magalhães T, Murphy K, Efremov A, Davidson TA, Molina-Navarro E, Wood KA, Ortegón-Aznar I (2023) How on Earth did that get there? Natural and human vectors of aquatic macrophyte global distribution. Hydrobiologia 850(7): 1515–1542. https://doi.org/10.1007/s10750-022-05107-0

Lobato-de Magalhães T, Murphy K, Tapia GJ, Davidson TA, Molina-Navarro E, de-Nova JA, Efremov A (2024) Global hotspots of endemicity, rarity and speciation of aquatic macrophytes. Marine and Freshwater Research 75: MF23121. https://doi.org/10.1071/MF23121

Lourenco M, Woodborne S (2023) Defining the Angolan Highlands Water Tower, a 40 plus-year precipitation budget of the headwater catchments of the Okavango Delta. Environmental Monitoring and Assessment 195(7): 859. https://doi.org/10.1007/s10661-023-11448-7

Lourenco M, Fitchett JM, Woodborne S (2022) Angolan highlands peatlands: Extent, age and growth dynamics. The Science of the Total Environment 810: 152315. https://doi.org/10.1016/j.scitotenv.2021.152315

Mahmoud WF, Farahat EA, Fahmy GM, Farrag HF, Awad HE (2021) Monthly and seasonal variations of biomass partitioning and macronutrients in the invasive grass Vossia cuspidata (Roxb.) Griff. Aquatic Botany 172: 103399. https://doi.org/10.1016/j.aquabot.2021.103399

Míguez C, Cancela Á, Sánchez Á, Álvarez X (2022) Possibilities for exploitation of invasive species, Arundo donax L., as a source of phenol compounds. Waste and Biomass Valorization 13(10): 4253–4265. https://doi.org/10.1007/s12649-022-01764-3

Milzow C, Kgotlhang L, Bauer-Gottwein P, Meier P, Kinzelbach W (2009) Regional review: The hydrology of the Okavango Delta, Botswana-processes, data and modelling. Hydrogeology Journal 17(6): 1297–1328. https://doi.org/10.1007/s10040-009-0436-0

Monteiro F, Costa E, Kissanga R, Costa JC, Catarino L (2022) An annotated checklist of the vascular flora of Quiçama National Park, Angola. Phytotaxa 557. https://doi.org/10.11646/phytotaxa.557.1.1

Moura-Júnior EGD, Cotarelli VM (2019) An update on the knowledge of aquatic macrophytes in Northeast Brazil. Rodriguésia 70: e04452017. https://doi.org/10.1590/2175-7860201970076

Moura-Júnior EGD, Lima LF, Silva SSL, Paiva RMS, Ferreira FA, Zickel CS, Pott A (2013) Aquatic macrophytes of Northeastern Brazil: Checklist, richness, distribution and life forms. Check List 9(2): 298–312. https://doi.org/10.15560/9.2.298

Moura-Júnior EGD, Paiva RMSD, Ferreira AC, Pacopahyba LD, Tavares AS, Ferreira FA, Pott A (2015) Updated checklist of aquatic macrophytes from Northern Brazil. Acta Amazonica 45(2): 111–132. https://doi.org/10.1590/1809-4392201402662

Murphy K, Efremov A, Davidson TA, Molina-Navarro E, Fidanza K, Betiol TCC, Urrutia-Estrada J (2019) World distribution, diversity and endemism of aquatic macrophytes. Aquatic Botany 158: 103127. https://doi.org/10.1016/j.aquabot.2019.06.006

Pestana MCA, Hora RC, Guarçoni EAE (2024) Floristic survey of aquatic macrophytes in eastern Maranhão, Brazil: Richness, biological forms and three new records. Brazilian Journal of Biology 84: e281276. https://doi.org/10.1590/1519-6984.281276

POWO (2024) Plants of the World Online. https://powo.science.kew.org/ [Acessed: 30 July 2024]

Ramsar (2025) The Convention on Wetlands. https://www.ramsar.org/news/angola-becomes-172nd-party-convention [Acessed: 01 May 2025]

Rejmánek M, Huntley BJ, Le Roux JJ, Richardson DM (2017) A rapid survey of the invasive plant species in western Angola. African Journal of Ecology 55(1): 56–69. https://doi.org/10.1111/aje.12315

Rohal CB, Cranney C, Hazelton EL, Kettenring KM (2019) Invasive Phragmites australis management outcomes and native plant recovery are context dependent. Ecology and Evolution 9(24): 13835–13849. https://doi.org/10.1002/ece3.5820

Roux JP (2009) Synopsis of the Lycopodiophyta and Pteridophyta of Africa, Madagascar and neighbouring islands. Strelitzia 23. South African National Biodiversity Institute, Pretoria.

Saleh MA (2023) Review of Ecological Significances and Roles of Aquatic Macrophytes. Fane-Fane International Multi-Disciplinary Journal 7: 458–468.

Shanungu GK (2009) Management of the invasive Mimosa pigra L. in Lochinvar National Park, Zambia. Biodiversity 10(2–3): 56–60. https://doi.org/10.1080/14888386.2009.9712844

Shelef O, Gross A, Rachmilevitch S (2013) Role of plants in a constructed wetland: Current and new perspectives. Water 5(2): 405–419. https://doi.org/10.3390/w5020405

Sieben E (2018) Inversodicraea warmingiana. [amended version of 2010 assessment] The IUCN Red List of Threatened Species 2018: e.T185590A124330759. https://doi.org/10.2305/IUCN.UK.2018-1.RLTS.T185590A124330759.en

Sieben E, Glen R, Muasya A (2017) The wetland flora of South Africa: Occurance patterns, freuquency and dominance across a diverse landscape. Aquatic Botany 142: 112–118. https://doi.org/10.1016/j.aquabot.2017.03.003

Sieben EJJ, Glen RP, Van Deventer H, Dayaram A (2021) The contribution of wetland flora to regional floristic diversity across a wide range of climatic conditions in southern Africa. Biodiversity and Conservation 30(3): 575–596. https://doi.org/10.1007/s10531-020-02104-4

Sood A, Uniyal PL, Prasanna R, Ahluwalia AS (2012) Phytoremediation potential of aquatic macrophyte, Azolla. Ambio 41: 122–137. https://doi.org/10.1007/s13280-011-0159-z

Sosef MS, Dauby G, Blach-Overgaard A, van Der Burgt X, Catarino L, Damen T, Couvreur TL (2017) Exploring the floristic diversity of tropical Africa. BMC Biology 15: 1–23. https://doi.org/10.1186/s12915-017-0356-8

Van Ginkel CE, Glen RP, Gordon-Gray KD, Cilliers CJ, Muasya M PP VD (2011) Easy identification of some South African wetland plants. Water Research Commission, Gezina.

Vymazal J (2013) Emergent plants used in free water surface constructed wetlands: A review. Ecological Engineering 61: 582–592. https://doi.org/10.1016/j.ecoleng.2013.06.023

Welgama A, Florentine S, Roberts J (2022) A global review of the woody invasive alien species Mimosa pigra (Giant Sensitive Plant): Its biology and management implications. Plants 11(18): 2366. https://doi.org/10.3390/plants11182366

Zigelski P, Gomes A, Finckh M (2019) Suffrutex dominated ecosystems in Angola. In: Huntley BJ, Russo V, Lages F, Ferrand N (Eds) Biodiversity of Angola. Science & conservation: a modern synthesis. Springer, 109–121. https://doi.org/10.1007/978-3-030-03083-4_7

Supplementary material 1

Botanical family, category of origin, distribution by province, life form, conservation status, citation and author

Authors: Rafael Somandjinga, Sílvia Quadros, Francisco Maiato, David Goyder

Data type: docx

Explanation note: A total of 526 aquatic macrophytes species of Angola, listed by their respective botanical family, category of origin, distribution by province, life form, conservation status, citation and author (voucher number).

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/phytokeys.259.147785.suppl1